



Theodore B. Shybut, M.D.
Sports Medicine & Arthroscopy
Department of Orthopedic Surgery
7200 Cambridge St. #10A
Houston, Texas 77030
Phone: 713-986-5590
Fax: 713-986-5521

GENERAL POST-OPERATIVE INSTRUCTIONS FOR SHOULDER SURGERY PATIENTS

The following information is general for all shoulder surgeries and may differ from your exact procedure.

• WOUND CARE & HYGIENE

General Shoulder Scopes (Requiring Only Small Portal Incisions)

Shoulder arthroscopies generally only require very small portal incisions to be made. Dr. Shybut usually allows you to remove the bulky dressing 72 hours after your surgery.

If you choose to remove your outer surgical dressing, please follow these guidelines. Remove the outer spongy compression tape along with the white gauze pads and white surgical padding. Bleeding usually occurs following surgery so these dressings may be soiled with blood. Only remove the elastic bandage and any white gauze pads or white surgical padding. Do NOT remove either the steri-strips that are directly on the skin or the yellow medicated gauze immediately on top of the steri-strips if at all possible. You are allowed to place water-proof bandages over the portal incisions to shower. Be sure to cover the entire area with water-proof bandages including the steri-strips and yellow gauze. Once you place water-proof bandages over your portals, leave these on until your first post-op appointment since removing these will remove the steri-strips and yellow gauze.

Open Procedures (Requiring Larger Incisions)

If you are having any type of procedure that requires a larger incision, Dr. Shybut does NOT allow you to remove your dressings after surgery. This includes you the patient, your caregivers, or the physical therapist. An exposed wound in physical therapy is UNACCEPTABLE. Removing your surgical dressings will expose you to potentially serious infections. Dressings will be changed on your first post-op visit and a new dressing applied. If your dressing gets EXCESSIVELY wet prior to your first post-op visit, meaning “soaked through,” please contact the office for instructions.

• POST-OPERATIVE SLING OR IMMOBILIZER – (IF APPLICABLE)

A post-operative brace with or without attached padded pillow may be necessary depending upon the specific procedure you will be having. This sling allows for limited to no movement of the shoulder in order to protect from unwanted motions following surgery. If a post-operative sling or immobilizer is ordered after surgery, Dr. Shybut will explain its use